

• 8.1.4

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \Gamma(\alpha, \beta) = \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}}$. Find the minimal sufficient statistics for:
 First note:

$$\frac{\Gamma(\vec{X} | \alpha, \beta)}{\Gamma(\vec{Y} | \alpha, \beta)} = \left(\frac{\prod x_i}{\prod y_i} \right)^\alpha e^{-\frac{\sum x_i - \sum y_i}{\beta}}$$

(a) and (b)

Clearly by the expression above, we have minimal sufficient statistics of $\prod x_i$ for α when β is known, and $\sum x_i$ for β when α is known.

(c) Show that MLE's in parts (a) and (b) are functions of the statistics in each part.
 MLE for α when β is known is the solution to

$$-n \ln(\beta) - \text{digamma}(\hat{\alpha}) + \sum \ln(x_i) = 0$$

this is clearly a function of our sufficient statistic from part (a) since $\sum \ln(x_i) = \ln(\prod x_i)$

MLE for β when α is known is

$$\frac{\bar{X}}{\alpha}$$

and this is clearly a function of the minimal sufficient statistic found in part (b).

Note: The method shown above is not unique for solving this problem. The Gamma distribution is a member of the exponential family (class) and thus we could have used this fact in solving for the sufficient statistics, and then could immediately have a confirmation on minimum sufficiency and completeness. However, the method above is not useless, not all distributions are exponential family, and then those results do not apply.

• 8.1.11

Let $X_1, \dots, X_n \stackrel{iid}{\sim} LN(\mu, \sigma^2)$. Find the joint sufficient statistics for (μ, σ^2) , and show that the MLE's are functions of this joint statistic.

We know already that for a r.v. $Y = \ln(X) \sim \mathcal{N}(\mu, \sigma^2)$, the sufficient statistic for $\theta = (\mu, \sigma^2)$ is $T(\vec{Y}) = (\sum y_i, \sum y_i^2)$. Thus our sufficient statistic for this problem will be $T(\vec{X}) = (\sum \ln(x_i), \sum (\ln(x_i))^2)$.

$$\hat{\mu} = \frac{\sum \ln(x_i)}{n} \quad (\text{fxn of sufficient statistic above})$$

$$\hat{\sigma}^2 = \frac{\sum \ln(x_i) - \hat{\mu}}{n} \quad (\text{also a fxn of sufficient statistic above})$$

• 8.1.12

Let $X_1, \dots, X_n \stackrel{iid}{\sim} f(x|\theta) = \theta(1+x)^{-(1+\theta)} 1_{(0,\infty)}(x)$ for $\theta > 0$. Show that $\prod_{i=1}^n (1+X_i)$ is a minimal sufficient statistic for θ , and that the MLE is a function of this statistic.

$$\frac{\prod_{i=1}^n \theta(1+x)^{-(1+\theta)} 1_{(0,\infty)}(x)}{\prod_{i=1}^n \theta(1+y)^{-(1+\theta)} 1_{(0,\infty)}(y)} = \frac{\theta^n (\prod_{i=1}^n (1+x_i))^{-(1+\theta)} 1_{(0,\infty)}(x_i)}{\theta^n (\prod_{i=1}^n (1+y_i))^{-(1+\theta)} 1_{(0,\infty)}(y_i)}$$

Now we can see that first, $\prod_{i=1}^n (1+x_i)$ is a sufficient by the 'factorization theorem', and second, that it is minimal since the expression will be independent of θ iff $\prod_{i=1}^n (1+x_i) = \prod_{i=1}^n (1+y_i)$.

Solving for the MLE for θ ,

$$\hat{\theta} = \frac{n}{\sum_{i=1}^n \ln(1+x_i)} = \frac{n}{\ln(\prod_{i=1}^n (1+x_i))}$$

and we can see this is a function of the minimal sufficient statistic found above. Note: "n" is in the numerator of our MLE, but that's okay! Sometimes it happens, as shown here.

• 8.2.4

(a) Let $X_1, X_2 \stackrel{iid}{\sim} f(x|\theta) = \frac{1}{\theta} e^{-\frac{x}{\theta}} 1_{(0,\infty)}(x)$. Show that $T_1(X_1, X_2) = X_1 + X_2$, and $T_2(X_1, X_2) = \frac{X_1}{X_1 + X_2}$ are independent r.v.'s. [Hint: Use Basu's Theorem (8.2.23)].

First thing to do is identify what distribution we're working with here, it's an exponential with mean parameter θ . To use Basu's theorem, we'll need to show two things: that one of T_1 , and T_2 is complete and sufficient, and the other is ancillary (not just first-order ancillary).

We already know $\sum_{i=1}^n x_i$ is sufficient for θ when we have $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{E}(\theta)$ from class. Thus, T_1 is sufficient here since we have $n = 2$. Now we just show T_2 is ancillary, that is, the distribution of T_2 is independent of θ .

First consider $Y_i = X_i/\theta$. A simple variable transformation shows $Y_i \sim \mathcal{E}(1)$ which is clearly independent of θ (see notes from 512, it's in there). Now note

$$T_2 = \frac{X_1}{X_1 + X_2} = \frac{\theta Y_1}{\theta(Y_1 + Y_2)} = \frac{Y_1}{Y_1 + Y_2}$$

Since the distribution of $Y_1/(Y_1 + Y_2)$ is independent of θ , and this equals T_2 , then the distribution of T_2 must also be independent of θ , and thus ancillary. Now by Basu's theorem, we have that T_1 , and T_2 are independent.

Slight aside: The exponential distribution (non-shifted one) belongs to a family called a "scale family". This is because they're all related by a scaled factor (as we saw above, by θ). There are also location families, and location-scale families. Similar to the method above for showing ancillarity, for location families we'll look for a difference between two r.v.'s, and for location-scale families we'll have both a difference and ratio (ex. $\frac{X_1 - X_2}{\sum X_i - X_2}$).

(b) Generalize the result in part (a) by proving:

If $X_1, \dots, X_n \stackrel{iid}{\sim} f(\text{part}(a))$, then $S_n = X_1 + \dots + X_n$ and $T_i = \frac{X_i}{S_n}$ are independent $\forall i \in \{1, \dots, n\}$.

Well we already said above that $S_n = \sum x_i$ is sufficient. Now we just note that instead of Y_1 , and Y_2 we now have Y_1, \dots, Y_n and we're done using the same argument(s) as before.

• 8.2.5

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{U}(0, \theta)$ and Y_i denote the i^{th} order statistic of the sample. Show that the distribution of $T_n = \sum_{i=1}^n X_i / Y_n$ does not depend on θ , hence that T_n is independent of Y_n .

One way to do this, as suggested by the hint, is through induction. But it kind of sucks, and there's an easier way.

First you have to note that the distribution we have here is a scale family as in the previous problem. Let $Z_i = \frac{X_i}{\theta}$

$$\begin{aligned} \frac{dx}{dz} &= \theta \\ f_Z(z) &= f_X(\theta \cdot z) \cdot \theta \\ &= \frac{\theta}{\theta} \mathbf{1}_{(0 \leq \theta \cdot z \leq \theta)} \\ &= \mathbf{1}_{(0 \leq z \leq 1)} \\ &= \mathcal{U}(0, 1) \\ T_n &= \sum_{i=1}^n \frac{X_i}{X_{(n)}} \\ &= \sum_{i=1}^n \frac{\theta Z_i}{\theta Z_{(n)}} \\ &= \sum_{i=1}^n \frac{Z_i}{Z_{(n)}} \end{aligned}$$

Since Z_i are independent of θ , and T_n can be expressed as only a function of the Z_i 's without θ , T_n must be an ancillary statistic. An example in the book showed that $X_{(n)}$ is a complete

sufficient statistic for θ , so by Basu's theorem, $Y_n = X_{(n)}$ and T_n are independent.

• 8.2.9

Let X_1, \dots, X_n be as in problem 8.1.8, that is, $\overset{iid}{\sim}$ discrete $\mathcal{U}(1, \theta)$. $Y_n = \max(X_1, \dots, X_n)$ is the minimal sufficient statistic for θ . Show Y_n is also complete.

$$\begin{aligned}
 P(Y_n \leq k) &= P(X \leq k)^n \\
 &= \frac{k^n}{\theta} \\
 P(Y_n = k) &= P(Y_n \leq k) - P(Y_n \leq k-1) \\
 &= \theta^{-n}(k^n - (k-1)^n) \\
 E(u(Y_n)) &= \sum_{k=1}^{\theta} u(k)\theta^{-n}(k^n - (k-1)^n) \\
 &= \theta^{-n} \sum_{k=1}^{\theta} u(k)(k^n - (k-1)^n) \\
 E(u(Y_n)) = 0 &\iff \sum_{k=1}^{\theta} u(k)(k^n - (k-1)^n) = 0
 \end{aligned}$$

We show $u(k)$ is identically zero by induction.

$$\begin{aligned}
 \text{Let } \theta &= 1 \\
 u(1) \cdot 1 &= 0 \\
 u(1) &= 0 \\
 \text{Assume } u(k) &= 0 \text{ for } k \in (1, \dots, \theta-1) \\
 \sum_{k=1}^{\theta} u(k)(k^n - (k-1)^n) &= u(\theta)(\theta^n - (\theta-1)^n) \\
 &= 0 \iff u(\theta) = 0
 \end{aligned}$$

Since $\theta^n - (\theta-1)^n \neq 0$ for $n > 0$. Therefore, Y_n is complete, since no function of it is an unbiased estimator of zero.

• 8.2.15

Let $X_1, \dots, X_n \overset{iid}{\sim}$ truncated Poisson where $p(x|\mu) = e^{-\mu}\mu^x/(x!(1 - e^{-\mu}))1_{\{1,2,\dots\}}(x)$.

(a) Show the joint distribution of X_1, \dots, X_n is a member of the exponential class.

$$\begin{aligned}
f(\vec{X} | \mu) &= \frac{e^{-n\mu} \mu^{\sum x_i}}{\prod x_i!} (1 - e^{-\mu})^{-n} = e^{\ln\left(\frac{e^{-n\mu} \mu^{\sum x_i}}{\prod x_i!} (1 - e^{-\mu})^{-n}\right)} \\
&= e^{\sum x_i \ln(\mu) - \ln(\prod x_i!) - n(\mu + \ln(1 - e^{-\mu}))}
\end{aligned}$$

Now we can see that we have the exponential family form where using the notation from D&M: $\phi_0(\vec{X}) = -\ln(\prod x_i!)$, $c_0(\mu) = -n(\mu + \ln(1 - e^{-\mu}))$, $\phi_1(\vec{X}) = \sum x_i$, $c_1(\mu) = \ln(\mu)$.

(b) Find a complete minimal sufficient statistic for μ .

From the form above and since our parameter space for μ , $\Theta = (0, \infty)$ contains a one-dimensional rectangle (interval), we can see that $\sum x_i$ is not only a sufficient statistic, but also minimal and complete.

• 8.2.17

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$. Show \bar{X}_n and $\sum_{i=1}^n a_i X_i$ are independent r.v.'s if $\sum_{i=1}^n a_i = 0$.

We already know that $\sum x_i$ is complete and minimally sufficient for μ when σ^2 is known (and hence \bar{X}_n). Now to use Basu's Theorem, we would just need to show that $\sum_{i=1}^n a_i X_i$ is ancillary with respect to μ .

Note that since each x_i is independently normally distributed, $\sum_{i=1}^n a_i X_i \sim \mathcal{N}(\sum_{i=1}^n a_i \mu, \sum_{i=1}^n a_i^2 \sigma^2) = \mathcal{N}(\mu \sum_{i=1}^n a_i, \sigma^2 \sum_{i=1}^n a_i^2)$. Now this will clearly be independent of μ (and thus ancillary) iff $\sum_{i=1}^n a_i = 0$. Thus by Basu's Theorem, \bar{X}_n and $\sum_{i=1}^n a_i X_i$ are independent r.v.'s if $\sum_{i=1}^n a_i = 0$.